cryptogenic cirrhosis

Cryptogenic cirrhosis is a condition that impairs liver function. People with this condition develop irreversible liver disease caused by scarring of the liver (cirrhosis), typically in mid- to late adulthood.

The liver is a part of the digestive system that helps break down food, store energy, and remove waste products, including toxins. Minor damage to the liver can be repaired by the body. However, severe or long-term damage can lead to the replacement of normal liver tissue with scar tissue.

In the early stages of cryptogenic cirrhosis, people often have no symptoms because the liver has enough normal tissue to function. Signs and symptoms become apparent as more of the liver is replaced by scar tissue. Affected individuals can experience fatigue, weakness, loss of appetite, weight loss, nausea, swelling (edema), enlarged blood vessels, and yellowing of the skin and whites of the eyes (jaundice).

People with cryptogenic cirrhosis may develop high blood pressure in the vein that supplies blood to the liver (portal hypertension). Cryptogenic cirrhosis can lead to type 2 diabetes, although the mechanism is unclear. Some people with cryptogenic cirrhosis develop cancer of the liver (hepatocellular cancer).

Frequency

Cirrhosis affects more than 600,000 people in the United States; cryptogenic cirrhosis likely accounts for 5 to 30 percent of these cases.

Genetic Changes

Unlike most cases of cirrhosis, cryptogenic cirrhosis is not caused by the hepatitis C or B virus or chronic alcohol use. A diagnosis of cryptogenic cirrhosis is typically given when all other causes of cirrhosis have been ruled out. When a disorder occurs without an apparent underlying reason, it is described as cryptogenic.

Research has shown that many cases of cryptogenic cirrhosis likely result from a condition called non-alcoholic fatty liver disease (NAFLD). In NAFLD, fat accumulates in the liver, impairing its function. If the fat buildup leads to inflammation and damage to liver tissue, NAFLD progresses to a condition called non-alcoholic steatohepatitis (NASH). Long term inflammation in people with NASH can cause the formation of scar tissue and a decrease in fat buildup. As a result, individuals progress from NASH to cirrhosis.

Cryptogenic cirrhosis may also develop from autoimmune hepatitis, which is a condition that occurs when the body's immune system malfunctions and attacks the liver, causing inflammation and liver damage.

In very rare cases, cryptogenic cirrhosis has been associated with mutations in genes that provide instructions for making certain keratin proteins. Keratins are a group of tough, fibrous proteins that form the structural framework of certain cells, particularly cells that make up the skin, hair, nails, and similar tissues. People with these keratin gene mutations are more likely to have fibrous deposits in their livers than individuals without the mutations. These deposits impair liver function, leading to cirrhosis. Mutations in these genes have also been found in people with other liver disorders.

In many cases, the cause of cryptogenic cirrhosis is unknown. Many people with predisposing conditions do not develop cirrhosis. Researchers are working to discover the causes of cryptogenic cirrhosis as well as to find out why some people seem to be protected from developing cirrhosis and others seem to be susceptible.

Inheritance Pattern

Most cases of cryptogenic cirrhosis are not inherited. However, people with a family history of liver disease or autoimmune disease are at an increased risk of developing these diseases themselves, and possibly cirrhosis.

In individuals with an associated keratin gene mutation, the risk of developing cryptogenic cirrhosis appears to have an autosomal dominant pattern of inheritance. Autosomal dominant inheritance means that one copy of an altered gene in each cell is sufficient to increase the risk of developing cryptogenic cirrhosis. In these families, people inherit an increased risk of cryptogenic cirrhosis, not the disease itself.

Other Names for This Condition

cirrhosis, cryptogenic

Diagnosis & Management

Genetic Testing

- Genetic Testing Registry: Cirrhosis, cryptogenic https://www.ncbi.nlm.nih.gov/gtr/conditions/C0267809/
- Genetic Testing Registry: Familial cirrhosis https://www.ncbi.nlm.nih.gov/gtr/conditions/C1861556/

Other Diagnosis and Management Resources

- Children's Hospital of Pittsburgh: Cirrhosis http://www.chp.edu/our-services/transplant/liver/education/liver-disease-states/cirrhosis
- Cleveland Clinic: Cirrhosis of the Liver http://my.clevelandclinic.org/health/articles/cirrhosis-of-the-liver
- MedlinePlus Encyclopedia: Cirrhosis https://medlineplus.gov/ency/article/000255.htm

General Information from MedlinePlus

- Diagnostic Tests
 https://medlineplus.gov/diagnostictests.html
- Drug Therapy https://medlineplus.gov/drugtherapy.html
- Genetic Counseling https://medlineplus.gov/geneticcounseling.html
- Palliative Care https://medlineplus.gov/palliativecare.html
- Surgery and Rehabilitation https://medlineplus.gov/surgeryandrehabilitation.html

Additional Information & Resources

MedlinePlus

- Encyclopedia: Cirrhosis https://medlineplus.gov/ency/article/000255.htm
- Encyclopedia: Fatty Liver--Nonalcoholic https://medlineplus.gov/ency/article/007657.htm
- Encyclopedia: Liver Cancer--Hepatocellular Carcinoma https://medlineplus.gov/ency/article/000280.htm
- Health Topic: Cirrhosis https://medlineplus.gov/cirrhosis.html

Additional NIH Resources

- National Cancer Institute: Liver (Hepatocellular) Cancer Prevention https://www.cancer.gov/types/liver/patient/liver-prevention-pdq#section/_21
- National Institute of Diabetes and Digestive and Kidney Diseases: Autoimmune Hepatitis https://www.niddk.nih.gov/health-information/liver-disease/autoimmune-hepatitis
- National Institute of Diabetes and Digestive and Kidney Diseases: Cirrhosis https://www.niddk.nih.gov/health-information/liver-disease/cirrhosis
- National Institute of Diabetes and Digestive and Kidney Diseases: Nonalcoholic Steatohepatitis
 https://www.niddk.nih.gov/health-information/liver-disease/nafld-nash

Educational Resources

- Disease InfoSearch: Cirrhosis of the Liver http://www.diseaseinfosearch.org/Cirrhosis+of+the+Liver/9662
- Johns Hopkins Medicine: Chronic Liver Disease/Cirrhosis
 http://www.hopkinsmedicine.org/healthlibrary/conditions/adult/liver_bili
 ary_and_pancreatic_disorders/chronic_liver_disease_cirrhosis_85,p00662/
- Johns Hopkins Medicine: Nonalcoholic Fatty Liver Disease http://www.hopkinsmedicine.org/healthlibrary/conditions/adult/digestive_disorders/nonalcoholic_fatty_liver_disease_134,178/
- MalaCards: cirrhosis, cryptogenic http://www.malacards.org/card/cirrhosis_cryptogenic
- Merck Manual Professional Version: Cirrhosis http://www.merckmanuals.com/professional/hepatic-and-biliary-disorders/fibrosis-and-cirrhosis/cirrhosis

Patient Support and Advocacy Resources

- American Liver Foundation: Cirrhosis
 http://www.liverfoundation.org/abouttheliver/info/cirrhosis/
- British Liver Trust: Cirrhosis https://www.britishlivertrust.org.uk/liver-information/liver-conditions/cirrhosis/
- Canadian Liver Foundation: Cirrhosis
 http://www.liver.ca/liver-disease/types/cirrhosis/default.aspx

ClinicalTrials.gov

ClinicalTrials.gov
 https://clinicaltrials.gov/ct2/results?cond=%22cryptogenic+cirrhosis%22

Scientific Articles on PubMed

PubMed

https://www.ncbi.nlm.nih.gov/pubmed?term=%28Liver+Cirrhosis%5BMAJR%5D%29+AND+%28cryptogenic+cirrhosis%5BTIAB%5D%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D

OMIM

 CIRRHOSIS, FAMILIAL http://omim.org/entry/215600

Sources for This Summary

- Caldwell S. Cryptogenic cirrhosis: what are we missing? Curr Gastroenterol Rep. 2010 Feb;12(1): 40-8. doi: 10.1007/s11894-009-0082-7. Review.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/20425483
- De BK, Mani S, Mandal SK, Mondal SS, Bhattacharya R, Pramanik AB, Sau D, Bhattacharjee K, Joardar S. Cryptogenic cirrhosis: metabolic liver disease due to insulin resistance. Indian J Med Sci. 2010 Nov;64(11):508-19. doi: 10.4103/0019-5359.102124.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/23051943
- Ku NO, Darling JM, Krams SM, Esquivel CO, Keeffe EB, Sibley RK, Lee YM, Wright TL, Omary MB. Keratin 8 and 18 mutations are risk factors for developing liver disease of multiple etiologies. Proc Natl Acad Sci U S A. 2003 May 13;100(10):6063-8. Epub 2003 Apr 30. Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/12724528
 Free article on PubMed Central: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC156326/
- Ku NO, Gish R, Wright TL, Omary MB. Keratin 8 mutations in patients with cryptogenic liver disease. N Engl J Med. 2001 May 24;344(21):1580-7.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/11372009
- Mohammed OK, Mahadeva S. Clinical outcomes of cryptogenic compared with non-cryptogenic cirrhosis: A retrospective cohort study. J Gastroenterol Hepatol. 2015 Sep;30(9):1423-8. doi: 10.1111/jgh.12978.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/25867030

Reprinted from Genetics Home Reference:

https://ghr.nlm.nih.gov/condition/cryptogenic-cirrhosis

Reviewed: March 2016 Published: March 21, 2017 Lister Hill National Center for Biomedical Communications U.S. National Library of Medicine National Institutes of Health Department of Health & Human Services